




Assembly Instructions for All Beetle Trap

IL-2700 – With Gel




- 1) Remove the clear top from the blue trap base by gently lifting from the anchor point.



- 2) Tear open and empty the contents of the pheromone gel into the tray.

Helpful tip: Squeeze the gel in a line around the perimeter of the tray to capture beetles as soon as they fall into the tray.



- 3) Snap the clear top onto the blue trap base.

Helpful tip: Tap the completed trap 2 to 3 times on a horizontal surface to settle the gel into the tray.

- 4) Place the trap on a flat surface where insect activity is suspected. Anchor the trap to equipment, walls or floor when necessary.











- 5) Inspect the trap regularly.



- 6) Replace the tray and pheromone gel every 60-90 days.

All Beetle Trap IL-2700 – With Gel

The SPB lure has pheromones and kairomones to attract the following beetles:

			
Red Flour Beetle (<i>Tribolium sp</i>)	Confused Flour Beetle (<i>Tribolium sp</i>)	Cigarette beetle (<i>Lasioderma serricorne</i>)	Warehouse beetle (<i>Trogoderma variabile</i>)
			
Rice weevil (<i>Sitophilus oryzae</i>)	Saw-toothed grain beetle (<i>Oryzaephilus spp</i>)	merchant grain beetle (<i>Oryzaephilus spp</i>)	Lesser grain borer (<i>Rhyzopertha dominica</i>)

Re-Order Information:

Photo	Component	Part Number
	PantryPatrol Pheromone Gel (Set of 10)	IL-1705-10
	All Beetle Trap Replacement Trays (Set of 10)	IL-1820-10